

Catatonia - Urgent Evaluation

Psychosis & Severe Mental Health

Catatonia - Urgent Evaluation is the focus of this educational guide. Psychotic symptoms can involve changes in perception, beliefs, thought organization, behavior, and the ability to judge what is happening. New, rapidly worsening, or dangerous symptoms require prompt in-person clinical assessment and sometimes emergency care. A symptom list cannot confirm a diagnosis: context, duration, severity, development, physical health, medication, substance use, and functional impact all matter.

Common patterns people may report

- marked reduction or increase in movement
- staring, mutism, posturing, or resistance to movement
- repetitive or unusual movements
- can occur in psychiatric or medical conditions and may require urgent in-person evaluation
- hearing, seeing, or sensing things that others do not
- fixed beliefs that remain strong despite contrary evidence
- marked suspiciousness or feeling watched or targeted
- speech or thinking that becomes difficult to follow

How it can affect everyday life

Depending on the person, these concerns may affect sleep, concentration, confidence, work or study performance, relationships, social activity, self-care, decision-making, or willingness to enter situations that feel uncertain. The level of distress and impairment is often more informative than any single symptom.

A useful first question

Instead of asking only 'Do I have this condition?', ask: What is happening, how often, for how long, in which situations, what does it stop me from doing, and what has already been tried? This produces information that is much more useful in a professional consultation.

What to notice before seeking help

- when perceptual or belief changes started
- sleep deprivation or substance use
- rapid change in functioning
- medication changes or medical illness
- ability to eat, drink, sleep, and care for yourself
- any risk of harm, severe agitation, or inability to stay safe

What else can look similar?

Psychological symptoms can overlap with sleep deprivation, medication effects, substance use, hormonal changes, pain, neurological or other medical conditions, and major life stress. A responsible assessment considers alternatives rather than assuming one explanation from an online checklist.

When professional assessment becomes especially important

- symptoms persist, intensify, or repeatedly return
- work, study, relationships, sleep, self-care, or daily functioning are significantly affected
- avoidance or coping behaviors are shrinking daily life
- there are safety concerns, self-harm thoughts, severe agitation, hallucinations, confusion, or inability to care for basic needs
- physical symptoms are new, severe, rapidly changing, or have not had appropriate medical evaluation

Urgent situations

SCUPsys consultations are not emergency services. If there is immediate danger, suicidal intent, severe confusion, rapidly worsening psychosis or mania, inability to care for basic needs, serious withdrawal symptoms, or a medical emergency, contact local emergency services or an appropriate in-person medical service now.

What a SCUPsys consultation can help with

A consultation can help organize the problem, identify patterns and triggers, clarify what information matters, provide psychoeducation, and outline sensible next steps. It can also help you decide what questions to take to a locally licensed clinician. It does not guarantee a diagnosis, prescription, medical certificate, or emergency treatment.

Prepare for a useful consultation

- Write a short timeline: when the concern began and what changed around that time.
- List the 3-5 symptoms or situations causing the most difficulty.
- Note sleep, stress, medication, substance use, recent illness, and major life events that may matter.
- Describe what you have already tried and what helped or made things worse.
- Choose one practical goal for the consultation: clarity, next-step planning, coping options, or deciding what type of local assessment is appropriate.

Use urgent in-person care for this presentation

This presentation can require immediate medical or psychiatric assessment. Do not delay urgent local care in order to purchase or wait for an online consultation. If immediate risk is present, contact local emergency services or go to an appropriate emergency medical service now. After the person is medically stabilized, SCUPsyc may be used only for non-urgent educational follow-up and planning.

Important boundaries

This guide and SCUPsyc's consultation services are educational and supportive. They are not a substitute for emergency care, a full diagnostic examination, medical testing, medication management, or treatment by a professional who is licensed and legally authorized to practice where the client is located. Availability and scope of remote services may vary by jurisdiction.

Authoritative starting points for further reading

World Health Organization: Mental disorders fact sheets and mental health resources

National Institute of Mental Health: Health Topics and public fact sheets

NHS: Mental health conditions and symptom guidance

Return to SCUPsyc Client Portal